arXiv:0908.4432v1 [math-ph] 30 Aug 2009

Superintegrability and higher order polynomial algebras II

Ian Marquette
Département de physique et Centre de recherche mathématique, Université de Montréal,
C.P.6128, Succursale Centre-Ville, Montréal, Québec H3C 3J7, Canada
ian.marquette@umontreal.ca

In an earlier article, we presented a method to obtain integrals of motion and polynomial algebras for a class of two-dimensional superintegrable systems from creation and annihilation operators. We discuss the general case and present its polynomial algebra. We will show how this polynomial algebra can be directly realized as a deformed oscillator algebra. This particular algebraic structure allows to find the unitary representations and the corresponding energy spectrum. We apply this construction to a family of caged anisotropic oscillators. The method can be used to generate new superintegrable systems with higher order integrals. We obtain new superintegrable systems involving the fourth Painlevé transcendent and present their integrals of motion and polynomial algebras.

1 Introduction

The following article is the second of a series of two [1] discussing the construction of higher order integrals of motion and polynomial algebras from creation and annihilation operators. We applied the results to two cases with a second and a third order integrals where no polynomial algebra were found

$$V(x,y)=\hbar^2[\frac{1}{8a^4}(x^2+y^2)+\frac{1}{y^2}+\frac{1}{(x+a)^2}+\frac{1}{(x-a)^2}], \tag{1.1}$$

$$V(x,y)=\hbar^2[\frac{1}{8a^4}(x^2+y^2)+\frac{1}{(y+a)^2}+\frac{1}{(y-a)^2}.+\frac{1}{(x+a)^2}+\frac{1}{(x-a)^2}]. \tag{1.2}$$

We constructed a quintic and a seventh order algebras. We studied the realization in terms of deformed oscillator algebras of a class of polynomial algebras of the seventh order. These results allowed us to obtain the structure function for the Potential 5 and 6 and unitary representations with the corresponding energy spectrum.

Many article [3-18] were devoted to superintegrable systems. However, most articles were on quadratically superintegrable systems. This paper follows also articles concerning superintegrable systems with third order integrals of motion [19,20,21,22,23,24]. The potentials given by Eq.(1.1) and (1.2) were obtained in Ref. 20 and studied from the point of view of supersymmetric quantum mechanics in Ref. 23. In Ref. 20 a potential written in terms of the fourth Painlevé transcendent was found.

$$V(x,y) = \frac{\omega^2}{2}(x^2+y^2) + \epsilon\frac{\hbar\omega}{2}f'(\sqrt{\frac{\omega}{\hbar}}x) + \frac{\omega\hbar}{2}f^2(\sqrt{\frac{\omega}{\hbar}}x) \tag{1.3}$$

$$+\omega\sqrt{\hbar\omega}xf(\sqrt{\frac{\omega}{\hbar}}x) + \frac{\hbar\omega}{3}(-\alpha+\epsilon) \quad .$$

We will use the potential given by Eq.(1.3) to generate new superintegrable systems with a higher order integrals of motion. By construction, they also have a second order integral of motion. This integral is related to separation of variables.

Let us present the organization of this paper. In Section 2, we recall how we can generate integrals of motion from the creation and annihilation operators. We consider the general case $m\lambda_x = n\lambda_y$ and obtain the polynomial algebra generated by integrals of motion. We show how we can obtain directly the realization in terms of deformed oscillator algebras. In Section 3, we apply the construction to a family of caged anisotropic harmonic oscillator. We show how the method can be used to generate new superintegrable systems with higher order integrals. We construct a new family of Hamiltonians written in terms of the fourth Painlevé transcendent. We present their integrals, polynomial algebra and energy spectrum. These results extend the number of known superintegrable systems involving the Painlevé transcendents

2 Polynomial algebras

Let us consider a two-dimension Hamiltonian separable in Cartesian coordinates

$$H(x,y,P_x,P_y) = H_x(x,P_x) + H_y(y,P_y), \tag{2.1}$$

for which creation and annihilation operators (polynomials in momenta) A_x, $A_x^\dagger$, A_y and $A_y^\dagger$ exist. These operators satisfy

$$[H_1, A_x^\dagger] = \lambda_x A_x^\dagger, \quad [H_2, A_y^\dagger] = \lambda_y A_y^\dagger \quad . \tag{2.2}$$

The following operators

$$f_1 = A_x^{\dagger m} A_y^n, \quad f_2 = A_x^m A_y^{\dagger n} \quad , \tag{2.3}$$

commute with the Hamiltonian H

$$[H, f_1] = [H, f_2] = 0, \tag{2.4}$$

if

$$m\lambda_x - n\lambda_y = 0, \quad m, n \in \mathbb{Z}^+ \quad . \tag{2.5}$$

Creation and annihilation operators allow to construct polynomial integrals of motion.
We will now consider integrals given by Eq.(2.7) and show that we can obtain directly a polynomial algebra written as a deformed oscillator algebra. Let us consider

$$H = H_x + H_y, \quad \lambda = m\lambda_x = n\lambda_y, \tag{2.6}$$

$$A = \frac{1}{2\lambda}(H_x - H_y), \quad I_- = A_x^m A_y^{\dagger n}, \quad I_+ = A_x^{\dagger m} A_y^n \tag{2.7}$$

We demand that creation and annihilation operators satisfy the following relations

$$[A_x, A_x^\dagger] = P(H_x) = Q(H_x + \lambda_x) - Q(H_x), \tag{2.8}$$

$$[A_y, A_y^\dagger] = R(H_y) = S(H_y + \lambda_y) - S(H_y). \tag{2.9}$$

We obtain the polynomial algebra

$$[A,I_-]=-I_-,\quad [A,I_+]=I_+, \tag{2.10}$$

$$[I_-,I_+]=\prod_{l=1}^{m}Q(\frac{H}{2}+m\lambda_x A+l\lambda_x)\prod_{k}^{n}S(\frac{H}{2}-n\lambda_y A-(n-k)\lambda_y) \tag{2.11}$$

$$-\prod_{i=1}^{m}Q(\frac{H}{2}+m\lambda_x A-(m-i)\lambda_x)\prod_{j}^{n}S(\frac{H}{2}-n\lambda_y A+j\lambda_y)$$

The order of the polynomial algebra is thus determine by the order of polynomials Q and S. This last relation have a very particular structure and we rewrite the Eq.(2.11) as

$$[I_-,I_+]=F_{m,n}(H,A+1)-F_{m,n}(H,A), \tag{2.12}$$

with

$$F_{m,n}=\prod_{i=1}^{m}Q(\frac{H}{2}+m\lambda_x A-(m-i)\lambda_x)\prod_{j}^{n}S(\frac{H}{2}-n\lambda_y A+j\lambda_y) \tag{2.13}$$

We can define

$$b^\dagger=I_+,\quad b=I_-,\quad N=A-u \tag{2.14}$$

$$[N,b^\dagger]=b^\dagger,\quad [N,b]=-b,\quad b^\dagger b=\Phi(H,N),\quad bb^\dagger=\Phi(H,N+1) \tag{2.15}$$

and

$$\Phi(H,N)=F_{m,n}(H,N+u) \tag{2.16}$$

To obtain Fock type unitary representations and their corresponding energy spectrum we impose the following conditions

$$\Phi(p+1,u_i,k)=0,\quad \Phi(0,u,k)=0,\quad \phi(x)>0,\quad \forall\quad x>0\quad . \tag{2.17}$$

We showed that the general polynomial algebra in the x and y axis given by Eq(2.2), (2.8) and (2.9) with $m\lambda_x=n\lambda_y$ allow us to generate a higher order polynomial algebra for the superintegrable system given by Eq.(2.10) and (2.11). This polynomial algebra is directly written as a deformed oscillator

algebra.
The two-dimensional anisotropic oscillator is a particular case and was studied in Ref. 25. From the point of view of physics this system is important in nuclear and atomic physics. In nuclear physics it describes single-particle level spectrum of pancake i.e. triaxially deformed nuclei with $\omega_x >> \omega_y, \omega_z$. The system with ratio 2 :1 describes superdeformed nuclei and ratio 3 :1 hyperdeformed nuclei. We will consider generalization of this systems that could have application in nuclear physics.

3 Applications

3.1 Caged anisotropic harmonic oscillator

The anisotropic harmonic oscillator can be generalized by adding singular terms. This system is the caged anisotropic harmonic oscillator [26,27]

$$H = \frac{P_x^2}{2} + \frac{P_y^2}{2} + \frac{\omega^2}{2}(k^2x^2 + m^2y^2) + \frac{l_1}{x^2} + \frac{l_2}{y^2}. \tag{3.1}$$

The method of separation of variables allows to solve the corresponding Schrödinger equation in terms of Laguerre polynomials [8] and to obtain the energy spectrum. However, the polynomial algebra remains to be determined. We apply to this system the construction of Section 2.

We have the following operators

$$\begin{aligned}
A_x^\dagger &= -\frac{1}{4}(\frac{\hbar}{\omega k}\frac{d^2}{dx^2} - 2x\frac{d}{dx} + \frac{\omega k}{\hbar}x^2 - \frac{2l_1}{\omega k \hbar x^2} - 1), \qquad (3.2)\\
A_x &= -\frac{1}{4}(\frac{\hbar}{\omega k}\frac{d^2}{dx^2} + 2x\frac{d}{dx} + \frac{\omega k}{\hbar}x^2 - \frac{2l_1}{\omega k \hbar x^2} + 1),\\
A_y &= -\frac{1}{4}(\frac{\hbar}{\omega m}\frac{d^2}{dy^2} - 2y\frac{d}{dy} + \frac{\omega m}{\hbar}y^2 - \frac{2l_2}{\omega m \hbar y^2} - 1),\\
A_y^\dagger &= -\frac{1}{4}(\frac{\hbar}{\omega m}\frac{d^2}{dy^2} + 2y\frac{d}{dy} + \frac{\omega m}{\hbar}y^2 - \frac{2l_2}{\omega m \hbar y^2} + 1).
\end{aligned}$$

They satisfy the relations given by Eq.(2.2) and (2.5) with $\lambda_x = 2\hbar k\omega$,

$\lambda_y = 2\hbar m\omega$ and $m\lambda_x = k\lambda_y$. We can apply results of Section 2. We have

$$Q(H_x) = \frac{1}{4\hbar^2 k^2\omega^2}H_x^2 - \frac{1}{2\hbar k\omega}H_x + (\frac{3}{16} - \frac{l_1}{2\hbar^2}), \tag{3.3}$$

$$S(H_y) = \frac{1}{4\hbar^2 m^2\omega^2}H_y^2 - \frac{1}{2\hbar m\omega}H_y + (\frac{3}{16} - \frac{l_2}{2\hbar^2}). \tag{3.4}$$

The integrals are given by Eq.(2.7). We have the condition $m, k \in \mathbb{Z}^+$. The Eq.(3.6) gives

$$\begin{aligned}\Phi_{m,k}(x) = m^2k^2\prod_{i=1}^{m}&(\frac{E}{4mk\hbar\omega} + x + u - 1 + \frac{i}{m} - \frac{1}{2m} - \frac{\nu_1}{2m})\\ &(\frac{E}{4mk\hbar\omega} + x + u - 1 + \frac{i}{m} - \frac{1}{2m} + \frac{\nu_1}{2m})\\ &\prod_{j=1}^{k}(\frac{E}{4mk\hbar\omega} - x - u + \frac{j}{k} - \frac{1}{2k} - \frac{\nu_2}{2k})\\ &(\frac{E}{4mk\hbar\omega} - x - u + \frac{j}{k} - \frac{1}{2k} + \frac{\nu_2}{2k}),\end{aligned} \tag{3.5}$$

with

$$\nu_1 = \sqrt{1 + \frac{8l_1}{\hbar^2}}, \quad \nu_2 = \sqrt{1 + \frac{8l_2}{\hbar^2}}.$$

We should impose the constraints given by the Eq.(2.17). We obtain the following solutions

$$u = \frac{-E}{4mk\hbar\omega} + \frac{m-p}{m} + \frac{1}{2m} + \frac{\epsilon_1\nu_1}{2m}, \tag{3.6}$$

$$\begin{aligned}\Phi_{m,k}(x) = m^2k^2\prod_{i=1}^{m}&(x + \frac{i-p}{m})(x + \frac{i-p}{m} + \frac{\epsilon_1\nu_1}{m})\\ \prod_{j=1}^{k}&(N + 1 + \frac{j-q}{k} - x)(N + 1 + \frac{j-q}{k} + \frac{\epsilon_2\nu_2}{k}),\end{aligned} \tag{3.7}$$

$$E = 2mk\hbar\omega(N + 2 + \frac{1 - 2p + \epsilon_1\nu_1}{2m} + \frac{1 - 2q + \epsilon_2\nu_2}{2k}). \tag{3.8}$$

$p = 1, 2, ..., m$, $q = 1, 2, ..., k$ and $N \in \mathbb{N}$.

3.2 System with Painlevé transcendent

In Ref. 20, five systems involving Painlevé transcendent [28] were found. One of these systems was written as a function of the fourth Painlevé transcendent.

$$V(x,y) = \frac{\omega^2}{2}(x^2+y^2) + \epsilon\frac{\hbar\omega}{2}f^{'}(\sqrt{\frac{\omega}{\hbar}}x) + \frac{\omega\hbar}{2}f^2(\sqrt{\frac{\omega}{\hbar}}x) \tag{3.9}$$
$$+\omega\sqrt{\hbar\omega}xf(\sqrt{\frac{\omega}{\hbar}}x) + \frac{\hbar\omega}{3}(-\alpha+\epsilon) \quad .$$

We presented its cubic algebra, wave functions and creation and annihilation operators. The Hamiltonians with $\epsilon = 1$ and $\epsilon = -1$ are related by a special case of third order supersymmetry called shape invariance [24,29].
Let us consider the following superintegrable system

$$H = \frac{P_x^2}{2} + \frac{P_y^2}{2} + g_1(x) + g_2(y) \quad , \tag{3.10}$$

$$g_1(x) = \frac{\omega_1^2}{2}x^2 + \frac{\hbar\omega_1\epsilon_1}{2}f_1^{'}(\sqrt{\frac{\omega_1}{\hbar}}x) + \frac{\omega_1\hbar}{2}f_1^2(\sqrt{\frac{\omega_1}{\hbar}}x) + \omega_1\sqrt{\hbar\omega_1}xf_1(\sqrt{\frac{\omega_1}{\hbar}}x) + \frac{\hbar\omega_1}{3}(-\alpha_1+\epsilon_1) \quad , \tag{3.11}$$

$$g_2(y) = \frac{\omega_2^2}{2}y^2 + \frac{\hbar\omega_2\epsilon_2}{2}f_2^{'}(\sqrt{\frac{\omega_2}{\hbar}}y) + \frac{\omega_2\hbar}{2}f_2^2(\sqrt{\frac{\omega_2}{\hbar}}y) + \omega_2\sqrt{\hbar\omega_2}yf_2(\sqrt{\frac{\omega_2}{\hbar}}y) + \frac{\hbar\omega_2}{3}(-\alpha_2+\epsilon_2) \quad , \tag{3.12}$$

with

$$m\omega_1 = n\omega_2, \quad \lambda_x = \hbar\omega_1 = \tilde{\omega}, \quad \lambda_y = \hbar\omega_2 \tag{3.13}$$

The function $f_1 = f_1(x,\alpha_1,\beta_1)$ and $f_2 = f_1(y,\alpha_2,\beta_2)$ are the fourth Painlevé transcendent. The third order creation and annihilation operators were discussed in Ref.24 and Ref.29. The commutator of these operators was also obtained. They satisfy the relation given by Eq(2.8) and (2.9) with

$$Q(H_x) = 8(H_x - \frac{\hbar\omega}{3}(-\alpha_1+\epsilon_1+3))((H_x - \frac{\hbar\omega}{3}(\frac{\alpha_1}{2}+4\epsilon_1-\frac{3}{2}))^2 + \frac{\omega^2\hbar^2\beta_1}{8}) \tag{3.14}$$

$$S(H_y) = 8(H_y - \frac{\hbar\omega}{3}(-\alpha_2+\epsilon_2+3))((H_y - \frac{\hbar\omega}{3}(\frac{\alpha_2}{2}+4\epsilon_2-\frac{3}{2}))^2 + \frac{\omega^2\hbar^2\beta_2}{8}) \tag{3.15}$$

The structure function of the general case is given by Eq.(2.16)

$$\Phi_{m,n}(x)=\prod_{i=1}^{m}(\frac{E}{2}+m\hbar\omega_1(x+u)-(m-i)\hbar\omega_1-\gamma_1 m\omega_1\hbar) \tag{3.16}$$

$$(\frac{E}{2}+m\hbar\omega_1(x+u)-(m-i)\hbar\omega_1-\gamma_2 m\omega_1\hbar)(\frac{E}{2}+m\hbar\omega_1(x+u)-(m-i)\hbar\omega_1-\gamma_3 m\omega_1\hbar)$$

$$\prod_{j=1}^{n}(\frac{E}{2}-n\hbar\omega_2(x+u)+j\hbar\omega_2-\gamma_4 n\omega_2\hbar)(\frac{E}{2}-n\hbar\omega_2(x+u)+j\hbar\omega_2-\gamma_5 n\omega_2\hbar)$$

$$(\frac{E}{2}-n\hbar\omega_2(x+u)+j\hbar\omega_2-\gamma_6 n\omega_2\hbar),$$

with

$$\gamma_1=-\frac{1}{3m}(-3+\alpha_1-\epsilon_1), \gamma_2=\frac{\hbar\omega_1}{12m}(-6+2\alpha_1-3i\sqrt{2\beta_1}+16\epsilon_1), \tag{3.17}$$

$$\gamma_3=\frac{\hbar\omega_1}{12m}(-6+2\alpha_1+3i\sqrt{2\beta_1}+16\epsilon_1), \gamma_4=\frac{1}{3n}(-3+\alpha_2-\epsilon_2), \tag{3.18}$$

$$\gamma_5=\frac{\hbar\omega_1}{12n}(-6+2\alpha_2-3i\sqrt{2\beta_2}+16\epsilon_2), \gamma_6=\frac{\hbar\omega_1}{12n}(-6+2\alpha_2+3i\sqrt{2\beta_2}+16\epsilon_2). \tag{3.19}$$

We can rewrite the structure function as

$$\Phi_{m,n}(x)=\prod_{i=1}^{m}\tilde{\omega}^6\hbar^6(\frac{E}{2\hbar\tilde{\omega}}+x+u-1+\frac{i}{m}-\gamma_1) \tag{3.20}$$

$$(\frac{E}{2\hbar\tilde{\omega}}+x+u-1+\frac{i}{m}-\gamma_2)(\frac{E}{2\hbar\tilde{\omega}}+x+u-1+\frac{i}{m}-\gamma_3)$$

$$\prod_{j=1}^{n}(\frac{E}{2\hbar\tilde{\omega}}-x-u+\frac{j}{n}-\gamma_4)(\frac{E}{2\hbar\tilde{\omega}}-x-u+\frac{j}{m}-\gamma_5)$$

$$(\frac{E}{2\hbar\tilde{\omega}}-x-u+\frac{j}{n}-\gamma_6).$$

We obtain the finite dimensional unitary representations and the corresponding energy spectrum from the Eq.(4.7). We have

$$u_1 = \frac{-E}{2\hbar\tilde{\omega}} + 1 - \frac{p}{m} + \gamma_1, u_2 = \frac{-E}{2\hbar\tilde{\omega}} + 1 - \frac{p}{m} + \gamma_2, u_3 = \frac{-E}{2\hbar\tilde{\omega}} + 1 - \frac{p}{m} + \gamma_3. \quad (3.21)$$

Let us present one of the three solutions for u_1

$$E = \hbar\tilde{\omega}(N + 2 - \frac{p}{m} - \frac{q}{n} + \gamma_1 + \gamma_4), \quad (3.22)$$

$$\Phi_1 = \prod_i^m \prod_j^n \tilde{\omega^6}\hbar^6 (x + \frac{i-p}{m})(x + \frac{i-p}{m} + \gamma_1 - \gamma_2) \quad (3.23)$$

$$(x + \frac{i-p}{m} + \gamma_1 - \gamma_3)(N + 1 - x + \frac{j-q}{n})$$

$$(N + 1 - x + \frac{j-q}{n} + \gamma_4 - \gamma_5)(N + 1 - x + \frac{j-q}{n} + \gamma_4 - \gamma_6).$$

$p = 1, 2, ..., m$, $q = 1, 2, ..., k$ and $N \in \mathbb{N}$.
There are 6 other solutions of the same form for u_2 and u_3.

4 Conclusion

The main result of this article is that we constructed in Section 2 the polynomial algebra for the general case $m\lambda = n\lambda$ and showed that this algebra can be realized directly as a deformed oscillator algebra. This result allows to study many systems that could have applications in nuclear physics, atomic physics and quantum chemistry.
We showed how the construction can be used to generate new superintegrable systems from known one-dimensional Hamiltonians with creation and annihilation operators. This result is also valid in classical mechanics.
We studied with this method a family of caged anisotropic oscillator. We found the polynomial algebra, the finite dimensional unitary representations and the energy spectrum. We applied also the method to a new superintegrable system involving the fourth Painlevé transcendent. We found the polynomial algebra, the finite dimensional unitary representations and the corresponding energy spectrum. The two systems given by Eq.(3.11), (3.10), (3.11) and (3.12) can be generalized in higher dimensions.

The classification of systems with creation and annhilation operators is important and could allow to find new superintegrable systems. A classification of second order creation and annihilation operators was discussed in Ref. 30. To our knowledge the classification of systems with second order ladder operators is not complete and only a class of Hamiltonians with third order ladder operators were discussed [29,24].

Acknowledgments The research of I.M. was supported by a postdoctoral research fellowship from FQRNT of Quebec. The author thanks P.Winternitz for very helpful comments and discussions.

5 References

1. I.Marquette, Superintegrability and higher order polynomial algebras I (2009).
2. V.A.Dulock and H.V.McIntosh, Am. J.Phys. 33, 109 (1965).
3. V.Fock, Z.Phys. 98, 145-154 (1935).
4. V.Bargmann, Z.Phys. 99, 576-582 (1936).
5. J.M.Jauch and E.L.Hill, Phys.Rev. 57, 641-645 (1940).
6. M.Moshinsky and Yu.F.Smirnov, The Harmonic Oscillator In Modern Physics, (Harwood, Amsterdam, 1966).
7. J.Fris, V.Mandrosov, Ya.A.Smorodinsky, M.Uhlir and P.Winternitz, Phys.Lett. 16, 354-356 (1965).
8. P.Winternitz, Ya.A.Smorodinsky, M.Uhlir and I.Fris, Yad.Fiz. 4, 625-635 (1966). (English translation in Sov. J.Nucl.Phys. 4, 444-450 (1967)).
9. A.Makarov, Kh. Valiev, Ya.A.Smorodinsky and P.Winternitz, Nuovo Cim. A52, 1061-1084 (1967).
10. N.W.Evans, Phys.Rev. A41, 5666-5676 (1990), J.Math.Phys. 32, 3369-3375 (1991).
11. E.G.Kalnins, J.M.Kress, W.Miller Jr and P.Winternitz, J.Math.Phys. 44(12) 5811-5848 (2003).
12. E.G.Kalnins, W.Miller Jr and G.S.Pogosyan, J.Math.Phys. A34, 4705-4720 (2001).
13. E.G.Kalnins, J.M.Kress and W.Miller Jr, J.Math.Phys. 46, 053509 (2005), 46, 053510 (2005), 46, 103507 (2005), 47, 043514 (2006), 47, 043514 (2006).
14. E.G.Kalnins, W.Miller Jr and G.S.Pogosyan, J.Math.Phys. 47, 033502.1-30 (2006), 48, 023503.1-20 (2007).

15. J.Daboul, P.Slodowy et C.Daboul, Phys.Lett. B317, 321-328 (1993)
16. Ya.I.Granovsky, A.S.Zhedanov et I.M.Lutzenko, Ann. Phys. (New York) 217, 1, 1-20 (1992).
17. P.Létourneau et L.Vinet, Ann. Phys. (New York) 243, 144-168 (1995).
18. C.Daskaloyannis, J.Math. Phys. 42, 1100-1119 (2001).
19. S.Gravel and P.Winternitz, J.Math.Phys. 43(12), 5902 (2002).
20. S.Gravel, J.Math.Phys. 45(3), 1003-1019 (2004).
21. I.Marquette and P.Winternitz, J.Math.Phys. 48(1) 012902 (2007).
22. I.Marquette and P.Winternitz, J. Phys. A : Math. Theor. 41, 304031 (2008).
23. I.Marquette, J. Math. Phys. 50, 012101 (2009).
24. I.Marquette, J.Math.Phys. 50 095202 (2009).
25. D.Bonatsos, C.Daskaloyannis, Prog.Part.Nucl.Phys. 43, 537 (1999).
26. P.E.Verrier and N.W.Evans, J.Math.Phys. 49, 022902 (2008).
27. M.A.Rodriguez, P.Tempesta and P.Wintertnitz, Phys. Rev. E78, 046608 (2008).
28. E.L.Ince, Ordynary Differential Equations, Dover, New York (1956).
29. A.Andrianov, F.Cannata, M.Ioffe and D.Nishnianidze, Phys.Lett.A, 266,341-349 (2000).
30. C.P.Boyer and W.Miller Jr., J.Math.Phys. 15, 9 (1974).